

CLICK HERE TO JUMP TO USE DIRECTIONS



TRIMEC[®] BENTGRASS FORMULA BROADLEAF HERBICIDE

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds.

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	9.92%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	6.12%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	2.53%
INERT INGREDIENTS:	81.43%
TOTAL 100.00%	

THIS PRODUCT CONTAINS:

- 0.71 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 8.20%.
 - 0.44 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.08%.
 - 0.18 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.10%.
- Isomer Specific by AOAC Methods.
Contains the single isomer form of Mecoprop-p.



KEEP OUT OF REACH OF CHILDREN

CAUTION

See below for additional Precautionary Statements and First Aid.

ONE GALLON COVERS UP TO 2.9 ACRES

KEEP FROM FREEZING



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. When mixing, loading, or applying this product, wear long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. <p style="text-align: right;"><i>(cont. on next column)</i></p>
--------------------	---

FIRST AID (continued)

If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.</p>	

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more in capacity: Do not open-pour from this container. A mechanical system (such as probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Environmental Hazards: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

For use on residential and other turf sites excluding sod farms.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solution near these plants. Do not spray on carpetgrass, dichondra nor lawns or turf where

desirable clovers are present. Use only lawn type sprayers. Coarse sprays are less likely to drift. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not spray while grass is emerging from dormancy. Do not apply to newly seeded grasses until well established. Do not spray when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application. Do not contaminate domestic or irrigation waters. Some injury can be expected to turf when used on centipedegrass or St. Augustinegrass lawns; only spot spray weeds on these types of grasses. Clover present will be damaged if sprayed. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

ORNAMENTAL LAWNS AND TURF

Bentgrass, Centipedegrass, St. Augustinegrass, and Buffalograss.

Apply Trimec® Bentgrass Formula Broadleaf Herbicide at 3 to 4 pints of product per acre (1.1-1.5 fl. oz. of product/1,000 sq. ft.) in sufficient water for uniform coverage. If hardy weeds persist, make a second application after three weeks. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. If application is made when soil is quite warm, use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until the second or third mowing. Seed can be sown 3 to 4 weeks after application. Do not use on dichondra. Do not apply this product to 'Floratam' St. Augustinegrass.

If dandelions persist, spot treatments with a single nozzle hand sprayer are recommended. Do not overdose bentgrass. In closely mowed bentgrass (putting and bowling greens) apply Trimec® Bentgrass Formula Broadleaf Herbicide at a maximum rate of 1 fl. oz. of product per 1,000 square feet preferably in May or through September. Any yellowing will disappear after about one week.

For hard-to-kill summer weeds, use 6 pints of product per acre. Do not apply this rate to bentgrass, centipedegrass or St. Augustinegrass.

The maximum application rate to turf is 0.4 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

NOTE: Care should be taken to avoid overdosing bentgrass or injury may result. High spray volumes will aid in obtaining uniform coverage. If hand operated sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom. Sweeping movements with a multiple nozzle boom will result in heavy over-application and subsequent turf discoloration or injury.

WEEDS CONTROLLED

Trimec® Bentgrass Formula Broadleaf Herbicide is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS and HENBIT. Other broadleaf weeds are listed below.

Aster, white heath & white prairie	False dandelion (*spotted catsear & common catsear)	Parsley-piert Pennsylvania smartweed
Bedstraw	Field bindweed	Pennywort (*dollarweed)
Beggarweed, creeping	(*morningglory & creeping jenny)	Pepperweed
Bindweed	Field oxeye-daisy	Pigweed
Black medic	(*creeping oxeye)	Pineappleweed
Broadleaf plantain	Filaree, whitestem & redstem	Plantain
Buckhorn plantain	Florida pusley	Poison ivy
Bull thistle	Ground ivy	Poison oak
Burclover	Groundsel	Prostrate knotweed (*knotweed)
Burdock, common	Hawkweed	Puncturevine
Buttercup, creeping	Healall	Purslane
Carpetweed	Henbit	Ragweed
Chickweed, common	Lambsquarters	Red sorrel (*sheep sorrel)
Chicory	Lawn burweed	Shepherdspurse
Cinquefoil	Lespedeza, common	Spotted spurge
Clover	Mallow, common	Spurge
Compassplant	Matchweed	Thistle
Curly dock	Mouseear chickweed	Veronica (*corn speedwell)
Dandelion	Oxalis (*yellow woodsorrel & creeping woodsorrel)	Virginia buttonweed (cont. on next column)
Dayflower		
Deadnettle		
Dock		
Dogfennel		
English daisy		

WEEDS CONTROLLED (cont.)

White clover (*Dutch clover, honeysuckle clover, white trefoil & purplewort)	Wild garlic Wild geranium Wild lettuce Wild mustard Wild onion Wild strawberry	Wild violet Yarrow Yellow rocket
Wild carrot		*Synonyms

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.

This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

840/9-2007 AP070203
EPA REG. NO. 2217-529

Manufactured By

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101
www.pbigordon.com